

DFUL-1/02/97

Integrable dissipative structures in the gauge theory of gravity

*L. Martina*¹, *O.K. Pashaev*^{†* 2} and *G. Soliani*³

*Dipartimento di Fisica dell'Università and INFN - Sezione di Lecce
73100 Lecce, Italy*

*†) Joint Institute for Nuclear Research, 141980 Dubna, Russia*⁴

Abstract

The Jackiw-Teitelboim gauge formulation of the 1+1 dimensional gravity allows us to relate different gauge fixing conditions with integrable hierarchies of evolution equations. We show that the equations for the Zweibein fields can be written as a pair of time reversed evolution equations of the reaction-diffusion type, admitting dissipative solutions. The spectral parameter for the related Lax pair appears as the constant valued spin connection associated with the $SO(1,1)$ gauge symmetry. Spontaneous breaking of the non-compact symmetry and the irreversible evolution are discussed.

PACS numbers: 04.70.Dy, 11.15-q, 11.10.Lm

¹E-mail: martina@le.infn.it

²E-mail: pashaev@main1.jinr.dubna.su

³E-mail: soliani@le.infn.it

⁴ Permanent address

1

In the last few years it has been shown that the classical treatment of the topological field theories can be strictly related to completely integrable models, which realize special non-covariant gauge conditions. At the classical level the topological field theories have zero Lagrangians on the solution manifold and the corresponding equations of motion coincide with the zero field strenghts. In a geometrical language this fact corresponds to the zero curvature condition required for the integrability of certain affine connections (the so-called linear problems), associated with the completely integrable systems.

On the other hand, it has been developed a systematic procedure in order to associate a completely integrable model with a Chern-Simons (CS) gauge theory [1, 2, 3]. This procedure has been performed for a compact group, but most of the involved algebra can be reused in the non-compact case. However, one can expect that the resulting nonlinear models may display some interesting physical and mathematical features. Indeed, from one side the non-compact groups appear typically in gravity theories as global symmetry groups and local symmetries in their gauge formulations. Furthermore, one knows that unstable and even chaotic dynamics is related to the hyperbolic and non-compact geometrical structures [4]. In the present work we consider the model called “lineal gravity”, introduced by Jackiw [5] and Teitelboim [6], in the form of a $SL(2, \mathbf{R})$ gauge theory in $(1+1)$ dimensions of the BF type [7, 8, 9, 10]. This model is equivalent to the Liouville gravity, whose quantization has been discussed recently [11]. In spite of its simple physical content, we are considering this theory as a mathematical laboratory in order to describe explicitly the abovementioned embedding. Although the conformal symmetry is generically broken, the integrable character of some of the resulting equations may provide a more direct application of the quantum inverse scattering. Furthermore, we show that the equations for the Zweibein fields are given by the pair of time-reversed nonlinear equations as in the thermo-field theory [15]. In fact, for the particular gauge condition provided by the $O(2,1)$ Heisenberg model, a system of reaction-diffusion equations arises, which appears in the theory of chemical reactions and in dissipative structures.

From the point of view of a gauge field theory of gravitation, the main idea is that the general coordinate transformations are implemented by gauge ones. Thus, the gauge groups induce a metric on a base space. On the other hand, one can always build up a non-compact nonlinear σ -model associated with certain zero-curvature conditions, resorting to the so-called moving trihedrals [12, 13]. In the following we shall study the class of σ -models related to the $SL(2, \mathbf{R})$ group. The algebra of this group and the corresponding Killing metric is given in terms of the basis elements τ_i ($i = 0, 1, 2$) by the bilinear product

$$\tau_i \tau_j = h_{ij} + i c_{ijk} \tau_k, \quad (1)$$

where $(h_{ij}) = \text{diag}(-1, -1, 1)$ and $c_{ijk} = -\epsilon_{ijl} h_{lk}$. With any point of a given

base space $\mathcal{M}$ endowed with the non positively definite $SL(2, \mathbf{R})$ metric we associate an orthonormal frame by a local adjoint representation of the algebra, expressed by

$$g\tau_i g^{-1} = n_i^k \tau_k. \quad (2)$$

The triad of vector fields $\mathbf{n}_i(x) = (h_{ij}n_j^i)$ would satisfy the orthonormal conditions

$$(\mathbf{n}_i \mathbf{n}_j) = h_{ij}, \quad \mathbf{n}_i \wedge \mathbf{n}_j = c_{ijk} \mathbf{n}_k. \quad (3)$$

If g is a differentiable function on $\mathcal{M}$, the local frame $\{\mathbf{n}_i\}$ changes accordingly to the adjoint representation of the right-invariant chiral currents $J_\mu = g^{-1} \partial_\mu g$, that is

$$\partial_\mu \mathbf{n}_i = (J_\mu)_{ik}^{(ad)} \mathbf{n}_k, \quad (4)$$

where $J_\mu^{(ad)}$ are matrices in the adjoint representation. Equations (4) can be seen as a linear system for the triad $\{\mathbf{n}_i\}$. Its integrability is assured by the zero curvature condition, satisfied by the chiral currents J_μ over a simply connected base space. Now we assign to one of the vector fields, say $\mathbf{n}_0 = \mathbf{s}$, the special role of (pseudo)-spin variable. It has the 1-sheeted hyperboloid $(\mathbf{s}, \mathbf{s}) = -1$ as phase space. The vector fields $\mathbf{n}_1, \mathbf{n}_2$ describe the tangent plane of such a hyperboloid and can be locally rotated. In other words a residual local $SO(1, 1)$ symmetry survives, corresponding to local Lorentz transformations. In order to specify this aspect, let us introduce a new basis $\mathbf{n}_\pm = \mathbf{n}_1 \pm \mathbf{n}_2$ and the following parametrization of the chiral currents

$$J_\mu = \frac{i}{4} \tau_0 V_\mu + \begin{pmatrix} 0 & q_\mu^- \\ q_\mu^+ & 0 \end{pmatrix}, \quad (5)$$

where V_μ and $q_\mu^\pm$ are real quantities defined by

$$V_\mu = 2(\mathbf{n}_2, \partial_\mu \mathbf{n}_1), \quad q_\mu^\pm = \pm \frac{1}{2}(\mathbf{s}, \partial_\mu \mathbf{n}_\pm). \quad (6)$$

This parametrization realizes a $\mathbf{Z}_2$ - graduation of the gauge algebra with isotropy group $\mathbf{R}$. Then, the system (4) takes the form

$$\begin{aligned} D_\mu^\mp \mathbf{n}_\pm &= \mp 2q_\mu^\pm \mathbf{s}, \\ \partial_\mu \mathbf{s} &= q_\mu^+ \mathbf{n}_- - q_\mu^- \mathbf{n}_+, \end{aligned} \quad (7)$$

where $D_\mu^\pm \equiv \partial_\mu \pm (1/2)V_\mu$ is the covariant derivative. This form is invariant under the local $SO(1, 1)$ gauge transformations generated by an arbitrary real function α

$$\mathbf{s} \rightarrow \mathbf{s}, \quad \mathbf{n}_+ \rightarrow e^{+\alpha} \mathbf{n}_+, \quad \mathbf{n}_- \rightarrow e^{-\alpha} \mathbf{n}_-, \quad (8)$$

or equivalently

$$V_\mu' = V_\mu + 2\partial_\mu \alpha, \quad q_\mu^{+'} = e^\alpha q_\mu^+, \quad q_\mu^{-'} = e^{-\alpha} q_\mu^-, \quad (9)$$

which are just the local boost rotations in the tangent plane to the hyperboloid described by $\mathbf{s}$. In this formalism, the zero curvature condition becomes

$$\begin{aligned} D_\mu^\mp q_\nu^\pm &= D_\nu^\mp q_\mu^\pm, \\ \partial_\mu V_\nu - \partial_\nu V_\mu &= 4(q_\mu^+ q_\nu^- - q_\nu^+ q_\mu^-). \end{aligned} \quad (10)$$

This system can be written in a base space of arbitrary dimension. However, it is natural to consider it in 2 or 3 dimensions. In the latter case Eq. (10) represents the Euler - Lagrange equations for the CS action on the $SL(2, \mathbf{R})$ group [14], which can be seen as a subgroup of the corresponding Poincaré group. On the other hand, in 2 - dimensions the local symmetry group identifies with the de Sitter group. In this framework one can introduce the action

$$\begin{aligned} S &= \frac{k}{2\pi} \int_{\mathcal{M}} Tr (J_0 F_{12}) dx^1 dx^2 \\ &= \frac{k}{2\pi} \int_{\mathcal{M}} \epsilon_{ab} \left[q_0^+ D_a^+ q_b^- + q_0^- D_a^- q_b^+ + \frac{1}{8} V^0 (\partial_a V_b - 4q_a^+ q_b^-) \right] dx^1 dx^2, \end{aligned} \quad (11)$$

where $F_{12} = \partial_1 J_2 - \partial_2 J_1 + [J_1, J_2]$, ϵ_{ab} is antisymmetric ($a, b = 1, 2$) and J_0 plays the role of Lagrangian multiplier. The field equations are $F_{12} = 0$ and $\mathcal{D}_a J_0 = 0$, where $\mathcal{D}_a = \partial_a - [J_a, \cdot]$. They are given equivalently by (10) for $\mu, \nu = 0, 1, 2$, under the assumption that all the derivatives of $q_\mu^\pm$ and V_μ with respect to x^0 vanish. This structure is a relict of the 3 - dimensional CS theory under the effects of the dimensional reduction. Of course, the space of all local solutions of the classical field equations, modulo gauge transformations, is finite dimensional. Furthermore, $\mathbf{s}(x^0, x^1, x^2)$ and $\mathbf{n}_\pm(x^0, x^1, x^2)$ can be found, by solving the linear differential equations (7) with x^0 -independent coefficients. In particular, for $\mu = 0$, the eigenvalues of this system are 0 and $\pm 2\sqrt{(J_0, J_0)}$. Remarkably, $\partial_\mu (J_0, J_0) = 0$. Thus, instead of the variables $\mathbf{s}$ and $\mathbf{n}_\pm$ one could use J_0 and $\partial_a J_0$ ($a = 1, 2$) as a local frame, treating in this framework the system in analogy with [16]. Action (11) allows us to take contact with the 2 - dimensional Jackiw - Teitelboim gravity model [5, 6], by introducing the metric tensor

$$g_{ab} = 4(q_a^+ q_b^- + q_b^+ q_a^-) = e_a^m e_b^n \eta_{mn} \quad , \quad (12)$$

where the Zweibein and the spin connection are given by

$$e_a^0 \pm e_a^1 = 2\sqrt{2}q_a^\pm, \quad \omega_a = \frac{1}{2}V_a \quad (13)$$

and the local flat metric is $\eta_{mn} = \text{diag}(-1, 1)$. In these settings Eq. (10) prescribes the constancy of the 2 - dimensional scalar curvature to the value of the cosmological constant, normalized to -1 , and the torsionless condition of the world-sheet manifold. Indeed, Equations (10) can be derived also from the action [5, 6]

$$S = \int_{\mathcal{M}} \sqrt{-g} V_0 (R + 1) dx^1 dx^2 \quad , \quad (14)$$

where V_0 is a world scalar Lagrange multiplier related to the dilaton field. Action (14) admits as local symmetry group the de Sitter group $SL(2, \mathbf{R})$, with local Lorentz generator $L = \frac{i}{2}\tau_0$. Furthermore, the gauge invariance of the theory based on (11) is equivalent to the general covariance of (14) on the solutions of Eqs. (10). Finally, the metric tensor g_{ab} has the direct and simple representation in terms of spin variables

$$g_{ab} = 2(\partial_a \mathbf{s}, \partial_b \mathbf{s}) \quad . \quad (15)$$

This formula enables us to give a gravitational interpretation of the σ -model we are going to introduce. In particular, Equation (15) expresses that the (pseudo)-spin $\mathbf{s}$ provides a special parametrization of the surface manifold.

2

The idea we follow is to add to (7) a differential constraint in the (x^1, x^2) space for $\mathbf{s}$, such that a completely integrable dynamics is introduced in order to partially fix the gauge freedom in a controlled fashion and allowing a residual local Lorentz covariance. Moreover, we concentrate on the "effective" space-time (x^1, x^2) , forgetting for the moment all what concerns the variable x^0 and the current J_0 . Precisely, we consider as a constraint the classical continuous Heisenberg model realized on the $SL(2, \mathbf{R})/SO(1, 1)$ coset space

$$\partial_2 \mathbf{s} = \mathbf{s} \wedge \partial_1^2 \mathbf{s} \quad . \quad (16)$$

Substitution into (7) leads

$$q_2^+ = D_1^- q_1^+, \quad q_2^- = -D_1^+ q_1^- . \quad (17)$$

These relations allow us to exclude the $q_2^\pm$ from the fields equations (10), obtaining

$$D_2^\mp q_1^\pm \mp (D_1^\mp)^2 q_1^\pm = 0, \quad (18)$$

$$\partial_2 V_1 - \partial_1 V_2 = 4\partial_1(q_1^+ q_1^-). \quad (19)$$

Defining the flat connection

$$A_2 = V_2 + 4(q_1^+ q_1^- - \rho), \quad A_1 = V_1, \quad (20)$$

where ρ is an arbitrary real constant, and gauging out it by a local $SO(1, 1)$ transformation $A_j = 2\partial_j \lambda$, $q^\pm = q_1^\pm e^{\pm \lambda}$ for any regular real function λ , we get the nonlinear reaction-diffusion system

$$\partial_2 q^\pm \mp \partial_1^2 q^\pm \pm 2(q^+ q^- - \rho)q^\pm = 0 \quad . \quad (21)$$

Here only the global $SO(1,1)$ $q^\pm \rightarrow e^{\pm\alpha} q^\pm$ invariance survives. The system (21) represents a particular form of a 2 - component reaction - diffusion system, playing an important role in synergetics [17, 18, 19].

The equations (21) come from the Lagrangian density

$$\mathcal{L} = \frac{1}{2}(q^+ \partial_2 q^- - q^- \partial_2 q^+) - \partial_1 q^+ \partial_1 q^- - (q^+ q^-)^2. \quad (22)$$

By resorting to the first order formalism [20] we obtain the canonical pairs

$$\{q^-(x), q^+(y)\} = \delta(x - y) \quad . \quad (23)$$

The Hamiltonian density is

$$H = \partial_1 q^+ \partial_1 q^- + (q^+ q^-)^2 \quad , \quad (24)$$

whose integral provides the total conserved energy. The conserved quantity associated with the hyperbolic scale invariance of the system (21) is $M = \int q^+(x) q^-(x) dx$. We will see below that M is interpreted as the total mass of a given configuration. It should be noticed that the mass integral is related to the energy of the Heisenberg model (16) by $\mathcal{E} = \int \partial_1 \mathbf{s} \partial_1 \mathbf{s} = 4M$. Furthermore, the corresponding momentum is $P = \int (q^+ \partial_1 q^- - q^- \partial_1 q^+) dx$. Actually, we will show that the system (21) is completely integrable and it admits infinitely many integrals in involution. This property comes from the relation with the model (16), for which we can write down the Lax pair representation. This phenomenon is well studied in the case of the compact $SU(2)/U(1)$ continuous Heisenberg model, where the correspondence is established with the nonlinear Schrödinger equation [21, 1].

In the gauge fixed by (16) the metric tensor (12) takes the form

$$\begin{aligned} g_{00} &= -8\partial_1 q^+ \partial_1 q^-, \\ g_{11} &= 8q^+ q^-, \\ g_{01} &= g_{10} = 4(q^- \partial_1 q^+ - q^+ \partial_1 q^-). \end{aligned} \quad (25)$$

From the comparison of these formulas with the conserved quantities shown above, we observe that the components g_{11} and g_{01} have the meaning of the mass and the momentum densities, respectively.

A simple symmetry analysis of Eq. (21) provides the time-inversion symmetry $q^+ \rightarrow q^-, q^- \rightarrow q^+, t \rightarrow -t$ and the space reflection symmetry $x \rightarrow -x$. By inspection one finds also the Galilei invariance $x \rightarrow x + 2vt, t \rightarrow t, q^\pm \rightarrow e^{\mp\alpha(x',t')} q^\pm(x',t')$, where $\alpha(x, t) = v^2 t + vx$.

However, the full symmetry structure of the system (21) is encoded into the Lax pair representation, which can be obtained by following the procedure adopted in [1]. The main idea is to reduce the space of the chiral currents (5) consistently with the constraints (17) and provide a grading, by exploiting

the freedom in the choice of the vector field V_μ . In fact, we first substitute the relations (17) and (20) into the expression (5) of J_a ($a = 1, 2$). The corresponding consistency condition provides a system of differential equations a little more general than (21), because of the explicit presence of A_a . However, we can gauge out these fields in terms of an arbitrary scalar function. At least locally, we can set $A_2 = -\frac{1}{2}A_1^2 = -8\zeta^2$, where ζ is a real constant. A further Galilei transformation furnishes the Lax pair for the system (21)

$$\begin{aligned} L_1 &= \partial_1 + \begin{pmatrix} \zeta & q^- \\ q^+ & -\zeta \end{pmatrix}, \\ L_2 &= \partial_2 + \begin{pmatrix} 2\zeta^2 - (q^+ q^- - \rho) & -(\partial_1 - 2\zeta)q^- \\ (\partial_1 + 2\zeta)q^+ & -2\zeta^2 + (q^+ q^- - \rho) \end{pmatrix}. \end{aligned} \quad (26)$$

Thus, we have associated the spectral parameter ζ of the Lax pair with a constant value of the $SO(1, 1)$ spin connection. The equations $L_1\Psi = 0$ and $L_2\Psi = 0$ provide the auxiliary linear problems, whose integrability is assured by (21). In the compact $U(1)$ gauge theory the analogous situation is interpreted as the value of the condensate state of the statistical gauge field [1]. The Lax pair (26) is of the Zakharov-Shabat type [22], where a rotation of $\pi/2$ in the complex plane of the spectral parameter is required. The Bäcklund transformations for the system (21) are

$$\partial_1 (q^\pm - \tilde{q}^\pm) = \sqrt{(q^+ - \tilde{q}^+) (q^- - \tilde{q}^-) - \mu} (q^\pm + \tilde{q}^\pm), \quad (27)$$

$$\begin{aligned} \partial_2 (q^\pm - \tilde{q}^\pm) &= \pm \sqrt{(q^+ - \tilde{q}^+) (q^- - \tilde{q}^-) - \mu} \partial_1 (q^\pm + \tilde{q}^\pm) \\ &\mp (q^+ q^- + \tilde{q}^+ \tilde{q}^-) (q^\pm - \tilde{q}^\pm). \end{aligned} \quad (28)$$

Thus, simple steady state solutions can be easily found. For example, for $\rho = 0$ and the initial trivial configuration $\tilde{q}^+ = 0 = \tilde{q}^-$ one finds:

a) for $\mu < 0$,

$$q^\pm(x, t) = \frac{a e^{\pm a^2 t}}{\sinh a(x - \xi)} \quad (|\mu| = a^2), \quad (29)$$

b) for $\mu > 0$,

$$q^\pm(x, t) = a \frac{e^{\mp a^2 t}}{\cos a(x - \xi)}, \quad (\mu = a^2) \quad (30)$$

with $x = x^1$ and $t = x^2$. For $a \rightarrow 0$ one obtains the simple rational solution $q^\pm = \frac{1}{x}$. In a similar way one can obtain the steady state periodic solutions, namely

$$q^\pm = b e^{\mp \frac{a}{2}(\xi - \xi_0)} \frac{sn[a(\xi - \xi_0); r]}{cn[a(\xi - \xi_0); r]}, \quad \text{or} \quad q^\pm = \frac{b e^{\mp \frac{a}{2}(\xi - \xi_0)}}{cn[a(\xi - \xi_0); s]}, \quad (31)$$

where the moduli $r = \frac{1}{|a|}\sqrt{a^2 - b^2}$ and $s = \frac{|a|}{\sqrt{a^2 + b^2}}$ are given in terms of the real arbitrary parameters a, b . These solutions present first order poles on the real axis, located at odd multiples of the complete elliptic integral of the first kind. Furthermore, the system (21) ($\rho = 0$) admits the analogous of the bright soliton solution for the Nonlinear Schrödinger Equation. In a moving frame coordinate $\xi = x - vt + \xi_0$, such a solution is given by

$$q^\pm(x, t) = \pm k e^{\pm[(k^2 - \frac{1}{4}v^2)t - \frac{1}{2}v(\xi - \xi_0)]} \text{sech } k\xi \quad . \quad (32)$$

It depends on the two real parameters k and v . But in contrast to the bright soliton, the above solution does not preserve the amplitude for the components $q^\pm$ independently. During the time evolution one of the fields is exponentially growing, while the other one is decaying. At the same time, the product $q^+ q^-$ has the usual solitonic shape. In the context of synergetics [17, 18, 19], where the dissipative structures are investigated, a growing mode is associated with the pattern formation. Consequently, for the solution (32) the name of "dissipaton" is suggested. In some sense it can be considered as the square root of the soliton. We note that the metric tensor is a regular bounded function for the dissipaton solution [27]. In the space of parameters (v, k) there exists the critical line $v = 2k$. For the solution (32) obtained with $v < 2k$, at the infinity one has $q^\pm \rightarrow 0$. At the critical value $v = 2k$ the solution is a steady state in the moving frame $q^\pm = \pm k e^{\pm k\xi_0} (1 \mp \tanh k\xi)$, with constant asymptotics $q^\pm \rightarrow \pm 2k e^{\pm k\xi_0}$ for $x \rightarrow \mp\infty$ and $q^\pm \rightarrow \pm 0$ for $x \rightarrow \pm\infty$. In the over-critical case $v > 2k$, we are led to $q^\pm \rightarrow \pm\infty$ for $x \rightarrow \mp\infty$ and $q^\pm \rightarrow \pm 0$ for $x \rightarrow \pm\infty$. Furthermore, by computing the first integrals $M = -2k$, $P = Mv$ and $E = -\frac{1}{4}Mv^2 - \frac{1}{12}M^3 = \frac{1}{2}Mv^2 - \frac{1}{12}M(9v^2 + M^2)$, for $k < 0$ one can naturally interpret M as the mass of a non-relativistic particle with momentum P and energy E . The second term in the expression of the energy describes the bound state energy arising from the nonlinear interaction. Actually, by standard techniques (see for instance [23]) one finds that the solution (32) is the first iteration of the N -soliton superposition formula

$$q^\pm[N] = 2\Delta^\pm[2N] / \Delta[2N], \quad (33)$$

where

$$\Delta[2N] = \begin{vmatrix} \psi_1^+ & \cdot & \cdot & \psi_{2N}^+ \\ \zeta_1^{N-1} \psi_1^+ & \cdot & \cdot & \zeta_{2N}^{N-1} \psi_{2N}^+ \\ \psi_1^- & \cdot & \cdot & \psi_{2N}^- \\ \zeta_1^{N-1} \psi_1^- & \cdot & \cdot & \zeta_{2N}^{N-1} \psi_{2N}^- \end{vmatrix}, \quad \Delta^\pm[2N] = \begin{vmatrix} \psi_1^\pm & \cdot & \cdot & \psi_{2N}^\pm \\ \zeta_1^N \psi_1^\pm & \cdot & \cdot & \zeta_{2N}^N \psi_{2N}^\pm \\ \psi_1^\mp & \cdot & \cdot & \psi_{2N}^\mp \\ \zeta_1^{N-2} \psi_1^\mp & \cdot & \cdot & \zeta_{2N}^{N-2} \psi_{2N}^\mp \end{vmatrix} \quad (34)$$

and $(\psi_k^+, \psi_k^-)^T$ are eigenfunctions of the auxiliary linear problem generated by (26) with $q^\pm = 0$ and spectral parameter ζ_k . The wavefunction for $q^\pm[N]$ is

given by $\Psi [N] = (1/\Delta [2N]) (\Delta^+ [2N+1], \Delta^- [2N+1])^T$, where

$$\Delta^\pm [2N+1] = \begin{vmatrix} \Delta^\pm [2N] & \zeta^i \psi^\pm \\ \zeta_k^{N-1} \psi_k^\mp & \zeta^{N-1} \psi^\mp \end{vmatrix}.$$

Then, one can build up a matrix eigenfunction $\Psi [N]$ belonging to $SL(2, \mathbf{R})$ and, by developping in the chosen basis of the algebra the quantity $\Psi^{-1} [N] \tau_0 \Psi [N]$, one finds solutions for the spin system (16). Finally, by using (25) (or (15)) on

For the model (21) we can write the analogue of the dark soliton, i.e.

$$q^\mp(x, t) = k e^{\mp[(\frac{1}{4}v^2 + 2k^2)t + \frac{1}{2}v(\xi - \xi_0)]} \tanh k\xi, \quad (35)$$

which becomes a time independent kink for $v^2 = 8k^2$. In contrast with the case (32), now it is impossible to choose the parameters in order to have vanishing boundary conditions at both infinities.

Finally, looking for homogeneous mass density solutions such that $q^+ q^- = \frac{1}{2} \chi(t)$ holds for any regular function $\chi(t)$, one finds

$$q^\pm(x, t) = \frac{c_\pm e^{\mp \chi_0}}{(t - t_0)^{\frac{1}{2} \pm 2c_+ c_-}} e^{\mp \frac{x^2}{4(t-t_0)}}, \quad (36)$$

where $c_\pm$, χ_0 and t_0 are arbitrary reals. This solution describes the nonlinear deformation of the Wiener solution for the heat equation with $q^+ q^- = \frac{c_+ c_-}{t - t_0}$.

In conclusion we would like to stress some points. First of all, the same scheme can be used in embedding into the Jackiw-Teitelboim model other completely integrable spin models, like the conformal invariant σ -model for the classical spin vector $\mathbf{s}$,

$$\partial_+ \partial_- \mathbf{s} - (\partial_+ \mathbf{s}, \partial_- \mathbf{s}) \mathbf{s} = 0 \quad . \quad (37)$$

After some manipulations on the abelian gauge field, also in this case one is led to the conformal invariant hyperbolic Sinh-Gordon equation

$$\partial_+ \partial_- \phi = 2(e^\phi - U_+ U_- e^{-\phi}), \quad (38)$$

where we have introduced the derivatives in the light-cone coordinates $\partial_\pm = \partial_2 \pm \partial_1$, $e^\phi = q_+^+ q_-^-$ and the arbitrary chiral functions $U_\pm = q_\pm^+ q_\pm^-$ (all $X_\pm^\pm$ quantities are given by analogous light-cone linear combinations). The tangent space representation of the gauge condition (37) is given by $D_-^+ q_+^+ = 0$, $D_-^+ q_+^- = 0$, in analogy with (17). Properly choosing $U_\pm$, one is lead to the Liouville equation or to conformally broken completely integrable equations, like the Sinh-Gordon and Cosh-Gordon equations. The model (37) is of its own interest and a detailed analysis has been performed, but it is out from the purposes of this work. Indeed, here we want to outline the appearance of dissipative structures

in gravity theory in the sense expressed by Eq. (21). This system is very similar to the "fictitious" or "mirror-image" systems with *negative* friction, which appear into the thermo-field approach to the damped harmonic oscillator treated in [15]. The energy is drained from the "real" oscillator to its "image", which mimics inaccessible states hidden in a thermostate. In this way the total energy is conserved and the Lagrangian description is allowed. That all this has some implication on certain thermodynamical properties of the gravity in the presence of black-holes [24, 25] is questionable, but suggestive.

Furthermore, inspired by the gauge equivalence between integrable models and related σ models, the idea of physical interpretation for gauge degrees of freedom can be useful. In fact, recently in [26] the analogy between point particles in 2+1 dimensional gravity and linear defects in solid continua was reconsidered in the framework of the CS gauge theory. In this analogy the essential difference arises from the role of diffeomorphisms which are gauge degrees of freedom (unobservable) in gravity, while in crystal physics they constitute elastic deformations, providing stress fields in the lattice.

Aknowledgments

This work was supported in part by MURST of Italy and by INFN - Sezione di Lecce. One of the authors (O. K. P.) thanks the Department of Physics of Lecce University for the warm hospitality and A. Fillipov, A. Pogrebkov and G. Vitiello for very useful discussions.

References

- [1] Pashaev O.K. 1996 *J Math Phys* **37** 4368
- [2] Martina L , Pashaev O K and Soliani G 1997 *J. Math. Phys.* **38** 1397
- [3] Martina L , Pashaev O K and Soliani G 1996 *Phys. Lett. B* **378** 175
- [4] Arnold V I , Avez A Ergodic Problems of Classical Mechanics 1968 W A Benjamin Inc. New York
- [5] Jackiw R 1984 in Quantum Theory of Gravity S. Christensen Ed. , Hilger, Bristol,
- [6] Teitelboim C 1983 *Phys. Lett. B* **126** 41
- [7] Isler K and Trugenberger C A 1989 *Phys Rev Lett* **63** 834
- [8] Chamseddine A H and Wyler D 1990 *Nucl Phys* **B340** 595
- [9] Montano D Sonnenschein J 1989 *Nucl. Phys.* **B324** 348
- [10] Birmingham D, Blau M, Rakowski M and Thompson G 1991 *Phys.Rep.* **209** 129
- [11] Seiberg N 1990 *Prog Theor Phys Suppl* **102** 319
- [12] Cartan E 1937 Théorie des Groupes Finis et Continus et la Géométrie Différentielle traitées par la Méthode du Repère Mobile, Paris, Gauthier-Villars
- [13] Orfanidis S 1980 *Phys.Rev.* **D21** 1513
- [14] Witten E (1988/89) *Nucl Phys* **B311** 46
- [15] Celeghini E, Rasetti M and Vitiello G 1992 *Ann. Phys.* **215** 156
- [16] Amati D, Elitzur S and Rabinovici E 1994 *Nucl. Phys.* **B418** 45
- [17] I Prigogine 1980 From being to becoming. Time and complexity in the physical sciences, W H Freeman and C., San Francisco
- [18] Haken H 1977 Synergetics Springer - Verlag, Berlin
- [19] Koch A J and Meinhardt H 1994 *Rev Mod Phys* **66** 1481
- [20] Faddeev L. and Jackiw R. 1988 *Phys. Rev. Lett.* **60**, 1692
- [21] Zakharov V E Takhtajan L 1979 *Theor. Math. Phys.* **38** 26
- [22] Zakharov V E Shabat A B 1971 *Zh. Eksp. Teor. Fiz.* **61** 118

- [23] Matveev V B and Salle M A 1991 *Darboux Transformations and Solitons*, Springer-Verlag, Berlin
- [24] Hawking S W 1975 *Comm. Math. Phys.* **43** 199
- [25] Israel W 1976 *Phys. Lett.* **A57** 107
- [26] Kohler C 1995 *Class. Quantum Grav.* **12** L11
- [27] However it admits causal singularities, corresponding to the horizons for black holes. Details of this interpretation would appear in a paper by the authors"